

STRUCTURAL RESULTS FOR FREE ARAKI-WOODS FACTORS AND THEIR CONTINUOUS CORES

CYRIL HOUDAYER

ABSTRACT. We show that for any type III_1 free Araki-Woods factor $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ associated with an orthogonal representation (U_t) of $\mathbf{R}$ on a separable real Hilbert space $H_{\mathbf{R}}$, the continuous core $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ is a semisolid II_{∞} factor, i.e. for any non-zero finite projection $q \in M$, the II_1 factor qMq is semisolid. If the representation (U_t) is moreover assumed to be mixing, then we prove that the core M is solid. As an application, we construct an example of a non-amenable solid II_1 factor N with full fundamental group, i.e. $\mathcal{F}(N) = \mathbf{R}_+^*$, which is not isomorphic to any interpolated free group factor $L(\mathbf{F}_t)$, for $1 < t \leq +\infty$.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

The *free Araki-Woods factors* were introduced by Shlyakhtenko in [32]. In the context of *free probability* theory, these factors can be regarded as the analogs of the *hyperfinit*e factors coming from the CAR^1 functor. To each real separable Hilbert space $H_{\mathbf{R}}$ together with an orthogonal representation (U_t) of $\mathbf{R}$ on $H_{\mathbf{R}}$, one can associate a von Neumann algebra denoted by $\Gamma(H_{\mathbf{R}}, U_t)''$, called the *free Araki-Woods von Neumann algebra*. The von Neumann algebra $\Gamma(H_{\mathbf{R}}, U_t)''$ comes equipped with a unique *free quasi-free state* denoted by φ_U , which is always normal and faithful on $\Gamma(H_{\mathbf{R}}, U_t)''$ (see Section 2 for a more detailed construction). If $\dim H_{\mathbf{R}} = 1$, then $\Gamma(\mathbf{R}, \text{Id})'' \cong L^{\infty}[0, 1]$. If $\dim H_{\mathbf{R}} \geq 2$, then $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ is a full factor. In particular, $\mathcal{M}$ can never be of type III_0 . The type classification of these factors is the following:

- (1) $\mathcal{M}$ is a type II_1 factor iff the representation (U_t) is trivial: in that case the functor Γ is Voiculescu's free Gaussian functor [38]. Then $\Gamma(H_{\mathbf{R}}, \text{Id})'' \cong L(\mathbf{F}_{\dim H_{\mathbf{R}}})$.
- (2) $\mathcal{M}$ is a type III_{λ} factor, for $0 < \lambda < 1$, iff the representation (U_t) is $\frac{2\pi}{|\log \lambda|}$ -periodic.
- (3) $\mathcal{M}$ is a type III_1 factor iff (U_t) is non-periodic and non-trivial.

Using free probability techniques, Shlyakhtenko obtained several remarkable classification results for $\Gamma(H_{\mathbf{R}}, U_t)''$. For instance, if the orthogonal representations (U_t) are *almost periodic*, then the free Araki-Woods factors $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ are completely classified up to state-preserving $*$ -isomorphism [32]: they only depend on Connes' invariant $\text{Sd}(\mathcal{M})$ which is equal in that case to the (countable) subgroup $S_U \subset \mathbf{R}_+^*$ generated by the eigenvalues of (U_t) . Moreover, the *discrete* core $\mathcal{M} \rtimes_{\sigma} \widehat{S_U}$ (where $\widehat{S_U}$ is the compact group dual of S_U) is $*$ -isomorphic to $L(\mathbf{F}_{\infty}) \otimes \mathbf{B}(\ell^2)$. Shlyakhtenko showed in [31] that if (U_t) is the left regular representation, then the *continuous* core $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ is isomorphic to $L(\mathbf{F}_{\infty}) \otimes \mathbf{B}(\ell^2)$ and the dual "trace-scaling" action (θ_s) is precisely the one constructed by Rădulescu [26]. For more on free Araki-Woods factors, we refer to [11, 13, 27, 28, 29, 30, 31, 32] and also to Vaes' Bourbaki seminar [37].

2000 *Mathematics Subject Classification.* 46L10; 46L54.

Key words and phrases. Free Araki-Woods factors; (Semi)solid factors; Deformation/Rigidity techniques; Intertwining techniques; Spectral gap property.

¹Canonical Anticommutation Relations

The free Araki-Woods factors as well as their continuous cores carry a *malleable deformation* in the sense of Popa. Then we will use the *deformation/rigidity* strategy together with the *intertwining techniques* in order to study the associated continuous cores. The high flexibility of this approach will allow us to work in a *semifinite* setting, so that we can obtain new structural/indecomposability results for the continuous cores of the free Araki-Woods factors. We first need to recall a few concepts. Following Ozawa [15, 16], a finite von Neumann algebra N is said to be:

- *solid* if for any diffuse von Neumann subalgebra $A \subset N$, the relative commutant $A' \cap N$ is amenable;
- *semisolid* if for any type II_1 von Neumann subalgebra $A \subset N$, the relative commutant $A' \cap N$ is amenable.

It is easy to check that solidity and semisolidity for II_1 factors are stable under taking amplification by any $t > 0$. Moreover, if N is a non-amenable II_1 factor, then $\text{solid} \implies \text{semisolid} \implies \text{prime}$. Recall in this respect that N is said to be *prime* if it cannot be written as the tensor product of two diffuse factors.

Ozawa discovered a class $\mathcal{S}$ of countable groups for which whenever $\Gamma \in \mathcal{S}$, the group von Neumann algebra $L(\Gamma)$ is solid [15]. He showed that the following countable groups belong to the class $\mathcal{S}$: the word-hyperbolic groups [15], the wreath products $\Lambda \wr \Gamma$ for Λ amenable and $\Gamma \in \mathcal{S}$ [16], and $\mathbf{Z}^2 \rtimes \text{SL}(2, \mathbf{Z})$ [17]. He moreover proved that if $\Gamma \in \mathcal{S}$, then for any free, ergodic, p.m.p. action $\Gamma \curvearrowright (X, \mu)$, the corresponding II_1 factor $L^\infty(X, \mu) \rtimes \Gamma$ is semisolid [16]. Recall that a non-amenable solid II_1 factor does not have property Γ of Murray & von Neumann [15].

Definition 1.1. Let M be a II_∞ factor and let Tr be a fixed faithful normal semifinite trace on M . We shall say that M is *solid* (resp. *semisolid*) if for any non-zero projection $q \in M$ such that $\text{Tr}(q) < \infty$, the II_1 factor qMq is solid (resp. semisolid).

Recall that an orthogonal/unitary representation (U_t) acting on H is said to be *mixing* if for any $\xi, \eta \in H$, $\langle U_t \xi, \eta \rangle \rightarrow 0$, as $|t| \rightarrow \infty$. The main result of this paper is the following:

Theorem 1.2. *Let $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ be a type III_1 free Araki-Woods factor. Then the continuous core $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ is a semisolid II_∞ factor. Since M is non-amenable, M is always a prime factor. If the representation (U_t) is moreover assumed to be mixing, then M is a solid II_∞ factor.*

The proof of Theorem 1.2 follows Popa's *deformation/rigidity* strategy. This theory has been successfully used over the last eight years to give a plethora of new classification/rigidity results for crossed products/free products von Neumann algebras. We refer to [5, 12, 14, 19, 20, 21, 22, 23, 24, 36] for some applications of the deformation/rigidity technique. We point out that in the present paper, the rigidity part does not rely on the notion of (relative) property (T) but rather on a certain *spectral gap* property discovered by Popa in [19, 20]. Using this powerful technique, Popa was able to show for instance that the Bernoulli action of groups of the form $\Gamma_1 \times \Gamma_2$, with Γ_1 non-amenable and Γ_2 infinite is $\mathcal{U}_{\text{fin}}$ -cocycle superrigid [19]. The spectral gap rigidity principle gave also a new approach to proving primeness and (semi)solidity for type II_1/III factors [4, 5, 19, 20]. We briefly remind below the concepts that we will play against each other in order to prove Theorem 1.2:

- (1) The first ingredient we will use is the “malleable deformation” by automorphisms (α_t, β) defined on the free Araki-Woods factor $\mathcal{M} * \mathcal{M} = \Gamma(H_{\mathbf{R}} \oplus H_{\mathbf{R}}, U_t \oplus U_t)''$. This deformation naturally arises as the “second quantization” of the rotations/reflection defined on $H_{\mathbf{R}} \oplus H_{\mathbf{R}}$ that commute with $U_t \oplus U_t$. It was shown in [19] that such a deformation automatically features a certain “transversality property” (see Lemma 2.1 in [19]) which will be of essential use in our proof.

- (2) The second ingredient we will use is the following property proved by Popa in [20], for free products of finite von Neumann algebras. Roughly, with B a finite amenable von Neumann algebra, any von Neumann subalgebra $Q \subset M$ with no amenable direct summand has “spectral gap” with respect to the orthogonal complement of M in $M *_B M$ (where M is regarded as the first copy in $M *_B M$). In other words, there exists a “critical” finite subset $F \subset \mathcal{U}(Q)$ such that if $x \in M *_B M$ almost commutes with all the unitaries $u \in F$, then x is almost contained in M .
- (3) A combination of (1) and (2) yields that for any $Q \subset M$ with no amenable direct summand, the malleable deformation (α_t) necessarily converges uniformly in $\|\cdot\|_2$ on $(Q' \cap M)_1$. Then, using the intertwining techniques, one can locate the position of $Q' \cap M$ inside M .

The second result of this paper provides a new example of a non-amenable solid II_1 factor. We first need the following:

Example 1.3. Using results of [2], we construct an example of an orthogonal representation (U_t) of $\mathbf{R}$ on a (separable) real Hilbert space $K_{\mathbf{R}}$ such that:

- (1) (U_t) is mixing.
- (2) The spectral measure of $\bigoplus_{n \geq 1} U_t^{\otimes n}$ is singular w.r.t. the Lebesgue measure on $\mathbf{R}$.

Shlyakhtenko showed in [31] that if the spectral measure of the representation $\bigoplus_{n \geq 1} U_t^{\otimes n}$ is singular w.r.t. the Lebesgue measure, then the continuous core of the free Araki-Woods factor $\Gamma(H_{\mathbf{R}}, U_t)''$ cannot be isomorphic to any $L(\mathbf{F}_t) \otimes \mathbf{B}(\ell^2)$, for $1 < t \leq \infty$, where $L(\mathbf{F}_t)$ denote the interpolated free group factors [8, 25]. Therefore, we obtain:

Theorem 1.4. *Let (U_t) be an orthogonal representation acting on $K_{\mathbf{R}}$ as in Example 1.3. Denote by $\mathcal{M} = \Gamma(K_{\mathbf{R}}, U_t)''$ the corresponding free Araki-Woods factor and by $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ its continuous core. Let $q \in L(\mathbf{R})$ be a non-zero projection such that $\text{Tr}(q) < \infty$. Then the non-amenable II_1 factor qMq is solid, has full fundamental group, i.e. $\mathcal{F}(qMq) = \mathbf{R}_+^*$, and is not isomorphic to any interpolated free group factor $L(\mathbf{F}_t)$, for $1 < t \leq \infty$.*

The paper is organized as follows. In Section 2, we recall the necessary background on free Araki-Woods factors as well as intertwining techniques for (semi)finite von Neumann algebras. Section 3 is mainly devoted to the proof of Theorem 1.2, following the deformation/spectral gap rigidity strategy presented above. In the last Section, we construct Example 1.3 and deduce Theorem 1.4.

Acknowledgement. The author would like to thank Prof. D. Shlyakhtenko for suggesting him the idea of looking at indecomposability results for the continuous cores of the free Araki-Woods factors and also for the useful discussions.

2. PRELIMINARY BACKGROUND

2.1. Shlyakhtenko’s free Araki-Woods factors. Let $H_{\mathbf{R}}$ be a real separable Hilbert space and let (U_t) be an orthogonal representation of $\mathbf{R}$ on $H_{\mathbf{R}}$ such that the map $t \mapsto U_t$ is strongly continuous. Let $H_{\mathbf{C}} = H_{\mathbf{R}} \otimes_{\mathbf{R}} \mathbf{C}$ be the complexified Hilbert space. We shall still denote by (U_t) the corresponding unitary representation of $\mathbf{R}$ on $H_{\mathbf{C}}$. Let A be the infinitesimal generator of (U_t) on $H_{\mathbf{C}}$ (Stone’s theorem), so that A is the positive, self-adjoint, (possibly) unbounded operator on $H_{\mathbf{C}}$ which satisfies $U_t = A^{it}$, for every $t \in \mathbf{R}$. Define another inner product on $H_{\mathbf{C}}$ by

$$\langle \xi, \eta \rangle_U = \left\langle \frac{2}{1 + A^{-1}} \xi, \eta \right\rangle, \forall \xi, \eta \in H_{\mathbf{C}}.$$

Note that for any $\xi \in H_{\mathbf{R}}$, $\|\xi\|_U = \|\xi\|$; also, for any $\xi, \eta \in H_{\mathbf{R}}$, $\Re(\langle \xi, \eta \rangle_U) = \langle \xi, \eta \rangle$, where $\Re$ denotes the real part. Denote by H the completion of $H_{\mathbf{C}}$ w.r.t. the new inner product

$\langle \cdot, \cdot \rangle_U$, and note that (U_t) is still a unitary representation on H . Introduce now the *full Fock space* of H :

$$\mathcal{F}(H) = \mathbf{C}\Omega \oplus \bigoplus_{n=1}^{\infty} H^{\otimes n}.$$

The unit vector Ω is called the *vacuum vector*. For any $\xi \in H$, we have the *left creation operator*

$$\ell(\xi) : \mathcal{F}(H) \rightarrow \mathcal{F}(H) : \begin{cases} \ell(\xi)\Omega = \xi, \\ \ell(\xi)(\xi_1 \otimes \cdots \otimes \xi_n) = \xi \otimes \xi_1 \otimes \cdots \otimes \xi_n. \end{cases}$$

For any $\xi \in H$, we denote by $s(\xi)$ the real part of $\ell(\xi)$ given by

$$s(\xi) = \frac{\ell(\xi) + \ell(\xi)^*}{2}.$$

The crucial result of Voiculescu [38] claims that the distribution of the operator $s(\xi)$ w.r.t. the vacuum vector state $\varphi_U = \langle \cdot, \Omega \rangle_U$ is the semicircular law of Wigner supported on the interval $[-\|\xi\|, \|\xi\|]$.

Definition 2.1 (Shlyakhtenko, [32]). Let (U_t) be an orthogonal representation of $\mathbf{R}$ on the real Hilbert space $H_{\mathbf{R}}$. The *free Araki-Woods* von Neumann algebra associated with $H_{\mathbf{R}}$ and (U_t) , denoted by $\Gamma(H_{\mathbf{R}}, U_t)''$, is defined by

$$\Gamma(H_{\mathbf{R}}, U_t)'' := \{s(\xi) : \xi \in H_{\mathbf{R}}\}''.$$

The vector state $\varphi_U = \langle \cdot, \Omega \rangle_U$ is called the *free quasi-free state*. It is normal and faithful on $\Gamma(H_{\mathbf{R}}, U_t)''$.

Recall that for any type III_1 factor $\mathcal{M}$, Connes-Takesaki's continuous decomposition [7, 33] yields

$$\mathcal{M} \otimes \mathbf{B}(L^2(\mathbf{R})) \cong (\mathcal{M} \rtimes_{\sigma} \mathbf{R}) \rtimes_{\theta} \mathbf{R},$$

where the *continuous core* $\mathcal{M} \rtimes_{\sigma} \mathbf{R}$ is a II_{∞} factor and θ is the *trace-scaling* action [33]:

$$\text{Tr}(\theta_s(x)) = e^{-s} \text{Tr}(x), \forall x \in (\mathcal{M} \rtimes_{\sigma} \mathbf{R})_+, \forall s \in \mathbf{R}.$$

The fact that $\mathcal{M} \rtimes_{\sigma} \mathbf{R}$ does not depend on the choice of a f.n. state on $\mathcal{M}$ follows from Connes' Radon-Nikodym derivative theorem [7]. Moreover, for any non-zero finite projection $q \in M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$, the II_1 factor qMq has full fundamental group.

Following [6], a factor $\mathcal{M}$ (with separable predual) is said to be *full* if the subgroup of inner automorphisms $\text{Inn}(\mathcal{M}) \subset \text{Aut}(\mathcal{M})$ is closed. Recall that $\text{Aut}(\mathcal{M})$ is endowed with the *u*-topology: for any sequence (θ_n) in $\text{Aut}(\mathcal{M})$,

$$\theta_n \rightarrow \text{Id}, \text{ as } n \rightarrow \infty \iff \|\varphi \circ \theta_n - \varphi\| \rightarrow 0, \text{ as } n \rightarrow \infty, \forall \varphi \in \mathcal{M}_*.$$

Since $\mathcal{M}$ has a separable predual, $\text{Aut}(\mathcal{M})$ is a polish group. For any II_1 factor N , N is full iff N does not have property Γ of Murray & von Neumann (see [6]).

Denote by $\pi : \text{Aut}(\mathcal{M}) \rightarrow \text{Out}(\mathcal{M})$ the canonical projection. Assume $\mathcal{M}$ is a full factor so that $\text{Out}(\mathcal{M})$ is a Hausdorff topological group. Fix a f.n. state φ on $\mathcal{M}$. Connes' invariant $\tau(\mathcal{M})$ is defined as the *weakest topology* on $\mathbf{R}$ that makes the map

$$\begin{aligned} \mathbf{R} &\rightarrow \text{Out}(\mathcal{M}) \\ t &\mapsto \pi(\sigma_t^{\varphi}) \end{aligned}$$

continuous. Note that this map does not depend on the choice of the f.n. state φ on $\mathcal{M}$ [7].

Denote by $\mathcal{F}(U_t) = \bigoplus_{n \in \mathbf{N}} U_t^{\otimes n}$. The modular group σ^{φ_U} of the free quasi-free state is given by: $\sigma_t^{\varphi_U} = \text{Ad}(\mathcal{F}(U_{-t}))$, for any $t \in \mathbf{R}$. The free Araki-Woods factors provided many new examples of full factors of type III [3, 7, 28]. We can summarize their general properties in the following theorem (see also Vaes' Bourbaki seminar [37]):

Theorem 2.2 (Shlyakhtenko, [28, 30, 31, 32]). *Let (U_t) be an orthogonal representation of $\mathbf{R}$ on the real Hilbert space $H_{\mathbf{R}}$ with $\dim H_{\mathbf{R}} \geq 2$. Denote by $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$.*

- (1) *$\mathcal{M}$ is a full factor and Connes' invariant $\tau(\mathcal{M})$ is the weakest topology on $\mathbf{R}$ that makes the map $t \mapsto U_t$ strongly continuous.*
- (2) *$\mathcal{M}$ is of type II_1 iff $U_t = \text{id}$ for every $t \in \mathbf{R}$. In this case, $\mathcal{M} \cong L(\mathbf{F}_{\dim(H_{\mathbf{R}})})$.*
- (3) *$\mathcal{M}$ is of type III_λ ($0 < \lambda < 1$) iff (U_t) is periodic of period $\frac{2\pi}{|\log \lambda|}$.*
- (4) *$\mathcal{M}$ is of type III_1 in the other cases.*
- (5) *$\mathcal{M}$ has almost periodic states iff (U_t) is almost periodic.*

Moreover, it follows from [27] that any free Araki-Woods factor $\mathcal{M}$ is *generalized solid* in the sense of [35]: for any diffuse von Neumann subalgebra $A \subset \mathcal{M}$ such that there exists a faithful normal conditional expectation $E : \mathcal{M} \rightarrow A$, the relative commutant $A' \cap \mathcal{M}$ is amenable.

Notice that the centralizer of the free quasi-free state $\mathcal{M}^{\varphi_U}$ may be trivial. This is the case for instance when the representation (U_t) has no eigenvectors. Nevertheless, the author recently proved in [11] that for any type III_1 free Araki-Woods factor $\mathcal{M}$, the *bicentralizer* is trivial, i.e. there always exists a faithful normal state ψ on $\mathcal{M}$ such that $(\mathcal{M}^\psi)' \cap \mathcal{M} = \mathbf{C}$. We refer to [10] for more on Connes' bicentralizer problem.

Remark 2.3 ([32]). Explicitly the value of φ_U on a word in $s(\xi_t)$ is given by

$$(1) \quad \varphi_U(s(\xi_1) \cdots s(\xi_n)) = 2^{-n} \sum_{(\{\beta_i, \gamma_i\}) \in \text{NC}(n), \beta_i < \gamma_i} \prod_{k=1}^{n/2} \langle \xi_{\beta_k}, \xi_{\gamma_k} \rangle_U.$$

for n even and is zero otherwise. Here $\text{NC}(2p)$ stands for all the non-crossing pairings of the set $\{1, \dots, 2p\}$, i.e. pairings for which whenever $a < b < c < d$, and a, c are in the same class, then b, d are not in the same class. The total number of such pairings is given by the p -th Catalan number

$$C_p = \frac{1}{p+1} \binom{2p}{p}.$$

Recall that a continuous φ -preserving action (σ_t) of $\mathbf{R}$ on a von Neumann algebra $\mathcal{M}$ endowed with a f.n. state φ is said to be *φ -mixing* if for any $x, y \in \mathcal{M}$ with $\varphi(x) = \varphi(y) = 0$,

$$(2) \quad \varphi(\sigma_t(x)y) \rightarrow 0, \text{ as } |t| \rightarrow \infty.$$

Proposition 2.4. *Let $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ be any free Araki-Woods factor and let φ_U be the free quasi-free state. Then*

$$(U_t) \text{ is mixing} \iff (\sigma_t^{\varphi_U}) \text{ is } \varphi_U\text{-mixing}.$$

Proof. We prove both directions.

$\Leftarrow$ For any $\xi, \eta \in H_{\mathbf{R}}$, $\varphi_U(s(\xi)) = \varphi_U(s(\eta)) = 0$. Moreover,

$$\begin{aligned} \langle U_t \xi, \eta \rangle_U &= 4\varphi_U(s(U_t \xi)s(\eta)) \\ &= 4\varphi_U(\sigma_{-t}^{\varphi_U}(s(\xi))s(\eta)) \rightarrow 0, \text{ as } |t| \rightarrow \infty. \end{aligned}$$

It follows that (U_t) is mixing.

$\Rightarrow$ Note that

$$\text{span} \{1, s(\xi_1) \cdots s(\xi_n) : n \geq 1, \xi_1, \dots, \xi_n \in H_{\mathbf{R}}\}$$

is a unital ultraweakly dense $*$ -subalgebra of $\mathcal{M}$. Using Kaplansky density theorem, it suffices to check Equation (2) for $x, y \in \mathcal{M}$ of the following form:

$$\begin{aligned} x &= s(\xi_1) \cdots s(\xi_{2k+1}) \\ y &= s(\eta_1) \cdots s(\eta_{2l+1}). \end{aligned}$$

Then

$$\begin{aligned} \varphi_U(\sigma_t^{\varphi_U}(x)y) &= \varphi(s(U_{-t}\xi_1) \cdots s(U_{-t}\xi_{2k+1})s(\eta_1) \cdots s(\eta_{2l+1})) \\ &= 2^{-2(k+l+1)} \sum_{(\{\beta_i, \gamma_i\}) \in \text{NC}(2(k+l+1)), \beta_i < \gamma_i} \prod_{j=1}^{k+l+1} \langle h_{\beta_j}, h_{\gamma_j} \rangle_U, \end{aligned}$$

where the letter h stands for $U_{-t}\xi$ or η . Notice that since $2k+1$ and $2l+1$ are odd, for any non-crossing pairing $(\{\beta_i, \gamma_i\}) \in \text{NC}(2(k+l+1))$, there must exist some $j \in \{1, \dots, k+l+1\}$ such that $\langle h_{\beta_j}, h_{\gamma_j} \rangle = \langle U_{-t}\xi_{\beta_j}, \eta_{\gamma_j} \rangle$. Since we assumed that (U_t) is mixing, it follows that $\varphi_U(\sigma_t^{\varphi_U}(x)y) \rightarrow 0$, as $|t| \rightarrow \infty$, and therefore $(\sigma_t^{\varphi_U})$ is φ_U -mixing. $\square$

Proposition 2.5. *Let $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$. If (U_t) is mixing, then Connes' invariant $\tau(\mathcal{M})$ is the usual topology on $\mathbf{R}$.*

Proof. Let $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$. Recall from Theorem 2.2 that $\tau(\mathcal{M})$ is the weakest topology on $\mathbf{R}$ that makes the map $t \mapsto U_t$ strongly continuous. Let (t_k) be a sequence in $\mathbf{R}$ such that $t_k \rightarrow 0$ w.r.t. the topology $\tau(\mathcal{M})$, as $k \rightarrow \infty$, i.e. $U_{t_k} \rightarrow \text{Id}$ strongly, as $k \rightarrow \infty$. Fix $\xi \in H_{\mathbf{R}}$, $\|\xi\| = 1$. Since

$$\lim_{k \rightarrow \infty} \langle U_{t_k} \xi, \xi \rangle = 1$$

and (U_t) is assumed to be mixing, it follows that (t_k) is necessarily bounded. Let $t \in \mathbf{R}$ be any cluster point for the sequence (t_k) . Then $U_t = \text{Id}$. Since (U_t) is mixing, it follows that $t = 0$. Therefore (t_k) converges to 0 w.r.t. the usual topology on $\mathbf{R}$. $\square$

2.2. Intertwining techniques for (semi)finite von Neumann algebras. Let (B, τ) be a finite von Neumann algebra with a distinguished f.n. trace. Since τ is fixed, we simply denote $L^2(B, \tau)$ by $L^2(B)$. Let H be a right Hilbert B -module, i.e. H is a complex (separable) Hilbert space together with a normal $*$ -representation $\pi : B^{\text{op}} \rightarrow \mathbf{B}(H)$. For any $b \in B$, and $\xi \in H$, we shall simply write $\pi(b^{\text{op}})\xi = \xi b$. By the general theory, we know that there exists an isometry $v : H \rightarrow \ell^2 \otimes L^2(B)$ such that $v(\xi b) = v(\xi)b$, for any $\xi \in H$, $b \in B$. Since $p = vv^*$ commutes with the right B -action on $\ell^2 \otimes L^2(B)$, it follows that $p \in \mathbf{B}(\ell^2) \otimes B$. Thus, as right B -modules, we have $H_B \simeq p(\ell^2 \otimes L^2(B))_B$.

On $\mathbf{B}(\ell^2) \otimes B$, we define the following f.n. semifinite trace Tr (which depends on τ): for any $x = [x_{ij}]_{i,j} \in (\mathbf{B}(\ell^2) \otimes B)_+$,

$$\text{Tr}([x_{ij}]_{i,j}) = \sum_i \tau(x_{ii}).$$

We set $\dim(H_B) = \text{Tr}(vv^*)$. Note that the dimension of H depends on τ but does not depend on the isometry v . Indeed take another isometry $w : H \rightarrow \ell^2 \otimes L^2(B)$, satisfying $w(\xi b) = w(\xi)b$, for any $\xi \in H$, $b \in B$. Note that $vw^* \in \mathbf{B}(\ell^2) \otimes B$ and $w^*w = v^*v = 1$. Thus, we have

$$\text{Tr}(vv^*) = \text{Tr}(vw^*vw^*) = \text{Tr}(wv^*vw^*) = \text{Tr}(ww^*).$$

Assume that $\dim(H_B) < \infty$. Then for any $\varepsilon > 0$, there exists a central projection $z \in \mathcal{Z}(B)$, with $\tau(z) \geq 1 - \varepsilon$, such that the right B -module H_z is finitely generated, i.e. of the form $pL^2(B)^{\oplus n}$ for some projection $p \in \mathbf{M}_n(\mathbf{C}) \otimes B$. The non-normalized trace on $\mathbf{M}_n(\mathbf{C})$ will be denoted by Tr_n .

In [21, 22], Popa introduced a powerful tool to prove the unitary conjugacy of two von Neumann subalgebras of a tracial von Neumann algebra (M, τ) . If $A, B \subset (M, \tau)$ are two (possibly non-unital) von Neumann subalgebras, denote by $1_A, 1_B$ the units of A and B . Note that we endow the finite von Neumann algebra B with the trace $\tau(1_B \cdot 1_B)/\tau(1_B)$.

Theorem 2.6 (Popa, [21, 22]). *Let $A, B \subset (M, \tau)$ be two (possibly non-unital) embeddings. The following are equivalent:*

- (1) There exist $n \geq 1$, a (possibly non-unital) $*$ -homomorphism $\psi : A \rightarrow B^n$ and a non-zero partial isometry $v \in \mathbf{M}_{1,n}(\mathbf{C}) \otimes 1_A M 1_B$ such that $xv = v\psi(x)$, for any $x \in A$.
- (2) The bimodule ${}_A L^2(1_A M 1_B)_B$ contains a non-zero sub-bimodule ${}_A H_B$ which satisfies $\dim(H_B) < \infty$.
- (3) There is no sequence of unitaries (u_k) in A such that $\|E_B(a^* u_k b)\|_2 \rightarrow 0$, as $k \rightarrow \infty$, for any $a, b \in 1_A M 1_B$.

If one of the previous equivalent conditions is satisfied, we shall say that A embeds into B inside M and denote $A \preceq_M B$.

Definition 2.7 (Popa & Vaes, [24]). Let $A \subset B \subset (N, \tau)$ be an inclusion of finite von Neumann algebras. We say that $B \subset N$ is *weakly mixing through A* if there exists a sequence of unitaries (u_k) in A such that

$$\|E_B(a^* u_k b)\|_2 \rightarrow 0, \text{ as } k \rightarrow \infty, \forall a, b \in N \ominus B.$$

The following result will be a crucial tool in Section 3: it will allow us to control the relative commutant $A' \cap N$ of certain subalgebras A of a given von Neumann algebra N .

Theorem 2.8 (Popa, [21]). Let (N, τ) be a finite von Neumann algebra and $A \subset B \subset N$ be von Neumann subalgebras. Assume that $B \subset N$ is weakly mixing through A . Then for any ${}_A H_B$ sub-bimodule of ${}_A L^2(N)_B$ such that $\dim(H_B) < \infty$, one has $H \subset L^2(B)$. In particular, $A' \cap N \subset B$.

For our purpose, we will need to use Popa's intertwining techniques for *semifinite* von Neumann algebras. We refer to Section 2 of [5] where such techniques were developed. Namely, let (M, Tr) be a von Neumann algebra endowed with a faithful normal semifinite trace Tr . We shall simply denote by $L^2(M)$ the M - M bimodule $L^2(M, \text{Tr})$, and by $\|\cdot\|_{2, \text{Tr}}$ the L^2 -norm associated with the trace Tr . We will use quite often the following inequality:

$$\|x\eta y\|_{2, \text{Tr}} \leq \|x\|_\infty \|y\|_\infty \|\eta\|_{2, \text{Tr}}, \forall \eta \in L^2(M), \forall x, y \in M,$$

where $\|\cdot\|_\infty$ denotes the operator norm. We shall say that a projection $p \in M$ is *Tr-finite* if $\text{Tr}(p) < \infty$. Note that a non-zero Tr-finite projection p is necessarily finite and $\text{Tr}(p \cdot p) / \text{Tr}(p)$ is a f.n. (finite) trace on pMp . Remind that for any projections $p, q \in M$, we have $p \vee q - p \sim q - p \wedge q$. Then it follows that for any Tr-finite projections $p, q \in M$, $p \vee q$ is still Tr-finite and $\text{Tr}(p \vee q) = \text{Tr}(p) + \text{Tr}(q) - \text{Tr}(p \wedge q)$.

Note that if a sequence (x_k) in M converges to 0 $*$ -strongly, as $k \rightarrow \infty$, then for any non-zero Tr-finite projection $q \in M$, $\|x_k q\|_{2, \text{Tr}} \rightarrow 0$, as $k \rightarrow \infty$. Indeed,

$$\begin{aligned} x_k \rightarrow 0 \text{ } * \text{-strongly in } M &\implies x_k^* x_k \rightarrow 0 \text{ weakly in } M \\ &\implies q x_k^* x_k q \rightarrow 0 \text{ weakly in } qMq \\ &\implies \text{Tr}(q x_k^* x_k q) \rightarrow 0 \\ &\implies \|x_k q\|_{2, \text{Tr}} \rightarrow 0. \end{aligned}$$

Moreover, there always exists an increasing sequence of Tr-finite projections (p_k) in M such that $p_k \rightarrow 1$ strongly, as $k \rightarrow \infty$.

Theorem 2.9 ([5]). Let (M, Tr) be a semifinite von Neumann algebra. Let $B \subset M$ be a von Neumann subalgebra such that $\text{Tr}|_B$ is still semifinite. Denote by $E_B : M \rightarrow B$ the unique Tr-preserving faithful normal conditional expectation. Let $q \in M$ be a non-zero Tr-finite projection. Let $A \subset qMq$ be a von Neumann subalgebra. The following conditions are equivalent:

- (1) There exists a Tr-finite projection $p \in B$, $p \neq 0$, such that the bimodule ${}_A L^2(qMp)_{pBp}$ contains a non-zero sub-bimodule ${}_A H_{pBp}$ which satisfies $\dim(H_{pBp}) < \infty$, where pBp is endowed with the finite trace $\text{Tr}(p \cdot p) / \text{Tr}(p)$.

- (2) *There is no sequence of unitaries (u_k) in A such that $E_B(x^* u_k y) \rightarrow 0$ $*$ -strongly, as $k \rightarrow \infty$, for any $x, y \in qM$.*

Definition 2.10. Under the assumptions of Theorem 2.9, if one of the equivalent conditions is satisfied, we shall still say that A *embeds into B inside M* and still denote $A \preceq_M B$.

3. STRUCTURAL RESULTS FOR THE CONTINUOUS CORES OF $\Gamma(H_{\mathbf{R}}, U_t)''$

3.1. Deformation/spectral gap rigidity strategy. We first introduce a few notations we will be using throughout this section. Let $H_{\mathbf{R}}$ be a separable real Hilbert space ($\dim(H_{\mathbf{R}}) \geq 2$) and let (U_t) be an orthogonal representation of $\mathbf{R}$ on $H_{\mathbf{R}}$ that we assume to be neither trivial nor periodic. We set:

- $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ is the free Araki-Woods factor associated with $(H_{\mathbf{R}}, U_t)$, φ is the free quasi-free state and σ is the modular group of the state φ . $\mathcal{M}$ is necessarily a type III₁ factor since (U_t) is neither periodic nor trivial.
- $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ is the continuous core of $\mathcal{M}$ and Tr is the semifinite trace associated with the state φ . M is a II_∞ factor since $\mathcal{M}$ is a type III₁ factor.
- Likewise $\widetilde{\mathcal{M}} = \Gamma(H_{\mathbf{R}} \oplus H_{\mathbf{R}}, U_t \oplus U_t)''$, $\widetilde{\varphi}$ is the corresponding free quasi-free state and $\widetilde{\sigma}$ is the modular group of $\widetilde{\varphi}$.
- $\widetilde{M} = \widetilde{\mathcal{M}} \rtimes_{\widetilde{\sigma}} \mathbf{R}$ is the continuous core of $\widetilde{\mathcal{M}}$ and $\widetilde{\text{Tr}}$ is the f.n. semifinite trace associated with $\widetilde{\varphi}$.

It follows from [32] that

$$(\widetilde{\mathcal{M}}, \widetilde{\varphi}) \cong (\mathcal{M}, \varphi) * (\mathcal{M}, \varphi).$$

In the latter free product, we shall write $\mathcal{M}_1$ for the first copy of $\mathcal{M}$ and $\mathcal{M}_2$ for the second copy of $\mathcal{M}$. We regard $\mathcal{M} \subset \widetilde{\mathcal{M}}$ via the identification of $\mathcal{M}$ with $\mathcal{M}_1$.

Denote by (λ_t) the unitaries in $L(\mathbf{R})$ that implement the modular action σ on $\mathcal{M}$ (resp. $\widetilde{\sigma}$ on $\widetilde{\mathcal{M}}$). Define the following faithful normal conditional expectations:

- $E : M \rightarrow L(\mathbf{R})$ such that $E(x\lambda_t) = \varphi(x)\lambda_t$, for every $x \in \mathcal{M}$ and $t \in \mathbf{R}$;
- $\widetilde{E} : \widetilde{M} \rightarrow L(\mathbf{R})$ such that $\widetilde{E}(x\lambda_t) = \widetilde{\varphi}(x)\lambda_t$, for every $x \in \widetilde{\mathcal{M}}$ and $t \in \mathbf{R}$.

Then

$$(\widetilde{M}, \widetilde{E}) \cong (M, E) *_{L(\mathbf{R})} (M, E).$$

Likewise, in the latter amalgamated free product, we shall write M_1 for the first copy of M and M_2 for the second copy of M . We regard $M \subset \widetilde{M}$ via the identification of M with M_1 . Notice that the conditional expectation E (resp. $\widetilde{E}$) preserves the canonical semifinite trace Tr (resp. $\widetilde{\text{Tr}}$) associated with the state φ (resp. $\widetilde{\varphi}$) (see [34]).

Consider the following orthogonal representation of $\mathbf{R}$ on $H_{\mathbf{R}} \oplus H_{\mathbf{R}}$:

$$V_s = \begin{pmatrix} \cos(\frac{\pi}{2}s) & -\sin(\frac{\pi}{2}s) \\ \sin(\frac{\pi}{2}s) & \cos(\frac{\pi}{2}s) \end{pmatrix}, \forall s \in \mathbf{R}.$$

Let (α_s) be the natural action on $(\widetilde{\mathcal{M}}, \widetilde{\varphi})$ associated with (V_s) : $\alpha_s = \text{Ad}(\mathcal{F}(V_s))$, for every $s \in \mathbf{R}$. In particular, we have

$$\alpha_s(s \begin{pmatrix} \xi \\ \eta \end{pmatrix}) = s(V_s \begin{pmatrix} \xi \\ \eta \end{pmatrix}), \forall s \in \mathbf{R}, \forall \xi, \eta \in H_{\mathbf{R}},$$

and the action (α_s) is $\widetilde{\varphi}$ -preserving. We can easily see that the representation (V_s) commutes with the representation $(U_t \oplus U_t)$. Consequently, (α_s) commutes with modular action $\widetilde{\sigma}$.

Moreover, $\alpha_1(x * 1) = 1 * x$, for every $a \in \mathcal{M}$. At last, consider the automorphism β defined on $(\widetilde{\mathcal{M}}, \widetilde{\varphi})$ by:

$$\beta(s \begin{pmatrix} \xi \\ \eta \end{pmatrix}) = s \begin{pmatrix} \xi \\ -\eta \end{pmatrix}, \forall \xi, \eta \in H_{\mathbf{R}}.$$

It is straightforward to check that β commutes with the modular action $\widetilde{\sigma}$, $\beta^2 = \text{Id}$, $\beta|_{\mathcal{M}} = \text{Id}|_{\mathcal{M}}$ and $\beta\alpha_s = \alpha_{-s}\beta$, $\forall s \in \mathbf{R}$. Since (α_s) and β commute with the modular action $\widetilde{\sigma}$, one may extend (α_s) and β to $\widetilde{M}$ by $\alpha_s|_{L(\mathbf{R})} = \text{Id}_{L(\mathbf{R})}$, for every $s \in \mathbf{R}$ and $\beta|_{L(\mathbf{R})} = \text{Id}_{L(\mathbf{R})}$. Moreover (α_s, β) preserves the semifinite trace $\widetilde{\text{Tr}}$. Let's summarize what we have done so far:

Proposition 3.1. *The $\widetilde{\text{Tr}}$ -preserving deformation (α_s, β) defined on $\widetilde{M}$ is s-malleable:*

- (1) $\alpha_s|_{L(\mathbf{R})} = \text{Id}_{L(\mathbf{R})}$, for every $s \in \mathbf{R}$ and $\alpha_1(x *_{L(\mathbf{R})} 1) = 1 *_{L(\mathbf{R})} x$, for every $x \in M$.
- (2) $\beta^2 = \text{Id}$ and $\beta|_M = \text{Id}|_M$.
- (3) $\beta\alpha_s = \alpha_{-s}\beta$, for every $s \in \mathbf{R}$.

Denote by $E_M : \widetilde{M} \rightarrow M$ the canonical trace-preserving conditional expectation. Since $\widetilde{\text{Tr}}|_M = \text{Tr}$, we will simply denote by Tr the semifinite trace on $\widetilde{M}$. Remind that the s-malleable deformation (α_s, β) automatically features a certain *transversality property*.

Proposition 3.2 (Popa, [19]). *We have the following:*

$$(3) \quad \|x - \alpha_{2s}(x)\|_{2, \text{Tr}} \leq 2 \|\alpha_s(x) - E_M(\alpha_s(x))\|_{2, \text{Tr}}, \forall x \in L^2(M, \text{Tr}), \forall s > 0.$$

The next proposition referred in the Introduction as the *spectral gap* property was first proved by Popa in [20] for free products of finite von Neumann algebras. We will need the following straightforward generalization:

Proposition 3.3 ([5]). *We keep the same notations as before. Let $q \in M$ be a non-zero projection such that $\text{Tr}(q) < \infty$. Let $Q \subset qMq$ be a von Neumann subalgebra with no amenable direct summand. Then for any free ultrafilter ω on $\mathbf{N}$, we have $Q' \cap (q\widetilde{M}q)^\omega \subset (qMq)^\omega$.*

Let $q \in M$ be a non-zero projection such that $\text{Tr}(q) < \infty$. Note that $\text{Tr}(q \cdot q)/\text{Tr}(q)$ is a finite trace on $q\widetilde{M}q$. If $Q \subset qMq$ has no amenable direct summand, then for any $\varepsilon > 0$, there exist $\delta > 0$ and a finite subset $F \subset \mathcal{U}(Q)$ such that for any $x \in (q\widetilde{M}q)_1$ (the unit ball w.r.t. the operator norm),

$$(4) \quad \|ux - xu\|_{2, \text{Tr}} < \delta, \forall u \in F \implies \|x - E_{qMq}(x)\|_{2, \text{Tr}} < \varepsilon.$$

We will simply denote $ux - xu$ by $[u, x]$.

3.2. Semisolidity of the continuous core. The following theorem is in some ways a reminiscence of a result of Ioana, Peterson & Popa, namely Theorem 4.3 of [14] and also Theorem 4.2 of [5]. The deformation/spectral gap rigidity strategy enables us to locate inside the core M of a free Araki-Woods factor the position of subalgebras $A \subset M$ with a *large* relative commutant $A' \cap M$.

Theorem 3.4. *Let $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ be a free Araki-Woods factor and $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ be its continuous core. Let $q \in L(\mathbf{R}) \subset M$ be a non-zero projection such that $\text{Tr}(q) < \infty$. Let $Q \subset qMq$ be a von Neumann subalgebra with no amenable direct summand. Then $Q' \cap qMq \preceq_M L(\mathbf{R})$.*

Corollary 3.5. *Let $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ be a free Araki-Woods factor of type III_1 . Then the continuous core $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ is a semisolid II_{∞} factor. Since M is non-amenable, M is always a prime factor.*

Proof of Theorem 3.4. Let $q \in L(\mathbf{R})$ be a non-zero projection such that $\text{Tr}(q) < \infty$. Let $Q \subset qMq$ be a von Neumann subalgebra with no amenable direct summand. Denote by $Q_0 = Q' \cap qMq$. We keep all the notations introduced previously and regard $M \subset \widetilde{M} = M_1 *_{L(\mathbf{R})} M_2$ via the identification of M with M_1 . Remind that $\alpha_s|_{L(\mathbf{R})} = \text{Id}_{L(\mathbf{R})}$, for every $s \in \mathbf{R}$. In particular $\alpha_s(q) = q$, for every $s \in \mathbf{R}$.

Step (1) : Using the spectral gap condition and the transversality property of (α_t, β) to find $t > 0$ and a non-zero intertwiner v between Id and α_t .

Let $\varepsilon = \frac{1}{4} \|q\|_{2, \text{Tr}}$. We know that there exist $\delta > 0$ and a finite subset $F \subset \mathcal{U}(Q)$, such that for every $x \in (q\widetilde{M}q)_1$,

$$\|[x, u]\|_{2, \text{Tr}} \leq \delta, \forall u \in F \implies \|x - E_{qMq}(x)\|_{2, \text{Tr}} \leq \varepsilon.$$

Since $\alpha_t \rightarrow \text{Id}$ pointwise $*$ -strongly, as $t \rightarrow 0$, and since F is a finite subset of $Q \subset qMq$, we may choose $t = 1/2^k$ small enough ($k \geq 1$) such that

$$\max\{\|u - \alpha_t(u)\|_{2, \text{Tr}} : u \in F\} \leq \frac{\delta}{2}.$$

For every $x \in (Q_0)_1$ and every $u \in F \subset Q$, since $[u, x] = 0$, we have

$$\begin{aligned} \|[\alpha_t(x), u]\|_{2, \text{Tr}} &= \|[\alpha_t(x), u - \alpha_t(u)]\|_{2, \text{Tr}} \\ &\leq 2\|u - \alpha_t(u)\|_{2, \text{Tr}} \\ &\leq \delta. \end{aligned}$$

Consequently, we get for every $x \in (Q_0)_1$, $\|\alpha_t(x) - E_{qMq}(\alpha_t(x))\|_{2, \text{Tr}} \leq \varepsilon$. Using Proposition 3.2, we obtain for every $x \in (Q_0)_1$

$$\|x - \alpha_s(x)\|_{2, \text{Tr}} \leq \frac{1}{2} \|q\|_{2, \text{Tr}},$$

where $s = 2t$. Thus, for every $u \in \mathcal{U}(Q_0)$, we have

$$\begin{aligned} \|u^* \alpha_s(u) - q\|_{2, \text{Tr}} &= \|u^*(\alpha_s(u) - u)\|_{2, \text{Tr}} \\ &\leq \|u - \alpha_s(u)\|_{2, \text{Tr}} \\ &\leq \frac{1}{2} \|q\|_{2, \text{Tr}}. \end{aligned}$$

Denote by $\mathcal{C} = \overline{\text{co}}^w\{u^* \alpha_s(u) : u \in \mathcal{U}(Q_0)\} \subset qL^2(\widetilde{M})q$ the ultraweak closure of the convex hull of all $u^* \alpha_s(u)$, where $u \in \mathcal{U}(Q_0)$. Denote by a the unique element in $\mathcal{C}$ of minimal $\|\cdot\|_{2, \text{Tr}}$ -norm. Since $\|a - q\|_{2, \text{Tr}} \leq 1/2 \|q\|_{2, \text{Tr}}$, necessarily $a \neq 0$. Fix $u \in \mathcal{U}(Q_0)$. Since $u^* a \alpha_s(u) \in \mathcal{C}$ and $\|u^* a \alpha_s(u)\|_{2, \text{Tr}} = \|a\|_{2, \text{Tr}}$, necessarily $u^* a \alpha_s(u) = a$. Taking $v = \text{pol}(a)$ the polar part of a , we have found a non-zero partial isometry $v \in q\widetilde{M}q$ such that

$$(5) \quad xv = v\alpha_s(x), \forall x \in Q_0.$$

Step (2) : Proving $Q_0 \preceq_M L(\mathbf{R})$ using the malleability of (α_t, β) . By contradiction, assume $Q_0 \not\preceq_M L(\mathbf{R})$. The first task is to lift Equation (5) to $s = 1$. Note that it is enough to find a non-zero partial isometry $w \in q\widetilde{M}q$ such that

$$xw = w\alpha_{2s}(x), \forall x \in Q_0.$$

Indeed, by induction we can go till $s = 1$ (because $s = 1/2^{k-1}$). Remind that $\beta(z) = z$, for every $z \in M$. Note that $vv^* \in Q'_0 \cap q\widetilde{M}q$. Since $Q_0 \not\preceq_M L(\mathbf{R})$, we know from Theorem 2.4 in

[5] that $Q'_0 \cap q\widetilde{M}q \subset qMq$. In particular, $vv^* \in qMq$. Set $w = \alpha_s(\beta(v^*)v)$. Then,

$$\begin{aligned} ww^* &= \alpha_s(\beta(v^*)vv^*\beta(v)) \\ &= \alpha_s(\beta(v^*)\beta(vv^*)\beta(v)) \\ &= \alpha_s\beta(v^*v) \neq 0. \end{aligned}$$

Hence, w is a non-zero partial isometry in $q\widetilde{M}q$. Moreover, for every $x \in Q_0$,

$$\begin{aligned} w\alpha_{2s}(x) &= \alpha_s(\beta(v^*)v\alpha_s(x)) \\ &= \alpha_s(\beta(v^*)xv) \\ &= \alpha_s(\beta(v^*x)v) \\ &= \alpha_s(\beta(\alpha_s(x)v^*)v) \\ &= \alpha_s\beta\alpha_s(x)\alpha_s(\beta(v^*)v) \\ &= \beta(x)w \\ &= xw. \end{aligned}$$

Since by induction, we can go till $s = 1$, we have found a non-zero partial isometry $v \in q\widetilde{M}q$ such that

$$(6) \quad xv = v\alpha_1(x), \forall x \in Q_0.$$

Note that $v^*v \in \alpha_1(Q_0)' \cap qMq$. Moreover, since $\alpha_1 : q\widetilde{M}q \rightarrow q\widetilde{M}q$ is a $*$ -automorphism, and $Q_0 \not\leq_M L(\mathbf{R})$, Theorem 2.4 in [5] gives

$$\begin{aligned} \alpha_1(Q_0)' \cap q\widetilde{M}q &= \alpha_1(Q'_0 \cap q\widetilde{M}q) \\ &\subset \alpha_1(qMq). \end{aligned}$$

Hence $v^*v \in \alpha_1(qMq)$.

Since $Q_0 \not\leq_M L(\mathbf{R})$, we know that there exists a sequence of unitaries (u_k) in Q_0 such that $E_{L(\mathbf{R})}(x^*u_ky) \rightarrow 0$ $*$ -strongly, as $k \rightarrow \infty$, for any $x, y \in qM$. We need to go further and prove the following:

Claim 3.6. $\forall a, b \in q\widetilde{M}q, \|E_{M_2}(a^*u_kb)\|_{2, \text{Tr}} \rightarrow 0$, as $k \rightarrow \infty$.

Proof of Claim 3.6. Let $a, b \in (\widetilde{M})_1$ be either elements in $L(\mathbf{R})$ or reduced words with letters alternating from $M_1 \ominus L(\mathbf{R})$ and $M_2 \ominus L(\mathbf{R})$. Write $b = yb'$ with

- $y = b$ if $b \in L(\mathbf{R})$;
- $y = 1$ if b is a reduced word beginning with a letter from $M_2 \ominus L(\mathbf{R})$;
- y = the first letter of b otherwise.

Note that either $b' = 1$ or b' is a reduced word beginning with a letter from $M_2 \ominus L(\mathbf{R})$. Likewise write $a = a'x$ with

- $x = a$ if $x \in L(\mathbf{R})$;
- $x = 1$ if a is a reduced word ending with a letter from $M_2 \ominus L(\mathbf{R})$;
- x = the last letter of a otherwise.

Either $a' = 1$ or a' is a reduced word ending with a letter from $M_2 \ominus L(\mathbf{R})$. For any $z \in Q_0 \subset M_1$, $xzy - E_{L(\mathbf{R})}(xzy) \in M_1 \ominus L(\mathbf{R})$, so that

$$E_{M_2}(azb) = E_{M_2}(a'E_{L(\mathbf{R})}(xzy)b').$$

Since $E_{L(\mathbf{R})}(xu_ky) \rightarrow 0$ strongly, as $k \rightarrow \infty$, it follows that $E_{M_2}(au_kb) \rightarrow 0$ strongly, as $k \rightarrow \infty$, as well. Thus, in the finite von Neumann algebra $q\widetilde{M}q$, we get $\|qE_{M_2}(au_kb)q\|_{2, \text{Tr}} \rightarrow 0$, as $k \rightarrow \infty$. Since $q \in L(\mathbf{R})$ and since the linear span of $L(\mathbf{R})$ and the reduced words with

letters alternating from $M_1 \ominus L(\mathbf{R})$ and $M_2 \ominus L(\mathbf{R})$ is a unital $*$ -strongly dense $*$ -subalgebra of $\widetilde{M}$, the rest of the proof follows from Kaplansky density theorem. $\square$

We remind that for any $x \in Q_0$, $v^*xv = \alpha_1(x)v^*v$. Moreover, $v^*v \in \alpha_1(qMq) \subset qM_2q$. So, for any $x \in Q_0$, $v^*xv \in qM_2q$. Since $\alpha_1(u_k) \in \mathcal{U}(qM_2q)$, we get

$$\|v^*v\|_{2,\text{Tr}} = \|\alpha_1(u_k)v^*v\|_{2,\text{Tr}} = \|E_{M_2}(\alpha_1(u_k)v^*v)\|_{2,\text{Tr}} = \|E_{M_2}(v^*u_kv)\|_{2,\text{Tr}} \rightarrow 0.$$

Thus $v = 0$, which is a contradiction. $\square$

Proof of Corollary 3.5. Let $q \in L(\mathbf{R})$ be a non-zero projection such that $\text{Tr}(q) < \infty$. Denote by $N = qMq$ the corresponding II_1 factor and by $\tau = \text{Tr}(q \cdot q)/\text{Tr}(q)$ the canonical trace on N . By contradiction, assume that N is not semisolid. Then there exists $Q \subset N$ a non-amenable von Neumann subalgebra such that the relative commutant $Q' \cap N$ is of type II_1 . Write $z \in \mathcal{Z}(Q)$ for the maximal projection such that Qz is amenable. Then $1 - z \neq 0$, the von Neumann algebra $Q(1 - z)$ has no amenable direct summand and $(Q' \cap N)(1 - z)$ is still of type II_1 . We may choose a projection $q_0 \in Q(1 - z)$ such that $\tau(q_0) = 1/n$. Since N is a II_1 factor, we may replace Q by $q_0Qq_0 \otimes \mathbf{M}_n(\mathbf{C})$, so that we may assume $Q \subset N$ has no amenable direct summand and $Q' \cap N$ is still of type II_1 .

If we apply Theorem 3.4, it follows that $Q' \cap N \preceq_M L(\mathbf{R})$. We get a contradiction because $Q' \cap N$ is of type II_1 and $L(\mathbf{R})$ is of type I. $\square$

It follows from [28] that for any type III_1 factor $\mathcal{M}$, if the continuous core $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ is full, then Connes' invariant $\tau(\mathcal{M})$ is the usual topology on $\mathbf{R}$. Let now $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ be a free Araki-Woods factor associated with (U_t) an almost periodic representation. Denote by $S_U \subset \mathbf{R}_+^*$ the (countable) subgroup generated by the point spectrum of (U_t) . Then $\tau(\mathcal{M})$ is strictly weaker than the usual topology. More precisely, the completion of $\mathbf{R}$ w.r.t. the topology $\tau(\mathcal{M})$ is the compact group $\widehat{S_U}$ dual of S_U (see [6]). Therefore in this case, for any non-zero projection $q \in L(\mathbf{R})$ such that $\text{Tr}(q) < \infty$, the II_1 factor qMq is semisolid, by Theorem 3.4, and has property Γ of Murray & von Neumann by the above remark.

3.3. Solidity of the continuous core under the assumption that (U_t) is mixing. We start this subsection with the following observations. The *solidity* of the continuous core M forces the centralizers on $\mathcal{M}$ to be *amenable*. Indeed, fix ψ any f.n. state on $\mathcal{M}$. Assume that the continuous core $M \simeq \mathcal{M} \rtimes_{\sigma^{\psi}} \mathbf{R}$ is solid. Choose a non-zero projection $q \in L(\mathbf{R})$ such that $\text{Tr}(q) < \infty$. Since $L(\mathbf{R})q$ is diffuse in $q(\mathcal{M} \rtimes_{\sigma^{\psi}} \mathbf{R})q$, its relative commutant must be amenable. In particular $\mathcal{M}^{\psi} \otimes L(\mathbf{R})q$ is amenable. Thus, $\mathcal{M}^{\psi}$ is amenable.

Note that if the orthogonal representation (U_t) contains a $\frac{2\pi}{|\log \lambda|}$ -periodic subrepresentation (V_t^{λ}) , $0 < \lambda < 1$, of the form

$$V_t^{\lambda} = \begin{pmatrix} \cos(t \log \lambda) & -\sin(t \log \lambda) \\ \sin(t \log \lambda) & \cos(t \log \lambda) \end{pmatrix},$$

then the free Araki-Woods factor $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ freely absorbs $L(\mathbf{F}_{\infty})$ (see [32]):

$$(\mathcal{M}, \varphi_U) * (L(\mathbf{F}_{\infty}), \tau) \cong (\mathcal{M}, \varphi_U).$$

In particular, the centralizer of the free quasi-free state $\mathcal{M}^{\varphi_U}$ is non-amenable since it contains $L(\mathbf{F}_{\infty})$. Therefore, whenever (U_t) contains a periodic subrepresentation of the form (V_t^{λ}) for some $0 < \lambda < 1$, the continuous core of $\Gamma(H_{\mathbf{R}}, U_t)''$ is semisolid by Theorem 3.4 but can never be solid. However, when (U_t) is assumed to be *mixing*, we get solidity of the continuous core. Indeed in that case, we can control the relative commutant $A' \cap M$ of diffuse subalgebras $A \subset L(\mathbf{R}) \subset M$, where M is the continuous core of the free Araki-Woods factor associated with (U_t) . Thus, the next theorem can be regarded as the analog of a result of Popa, namely Theorem 3.1 of [21] (see also Theorem D.4 in [36]).

Theorem 3.7. *Let (U_t) be a mixing orthogonal representation of $\mathbf{R}$ on the real Hilbert space $H_{\mathbf{R}}$. Denote by $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ the corresponding free Araki-Woods factor and by $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ its continuous core. Let $k \geq 1$ and let $q \in \mathbf{M}_k(\mathbf{C}) \otimes L(\mathbf{R})$ be a non-zero projection such that $T := (\text{Tr}_k \otimes \text{Tr})(q) < \infty$. Write $L(\mathbf{R})^T := q(\mathbf{M}_k(\mathbf{C}) \otimes L(\mathbf{R}))q$ and $M^T := q(\mathbf{M}_k(\mathbf{C}) \otimes M)q$. Let $A \subset L(\mathbf{R})^T$ be a diffuse von Neumann subalgebra.*

Then for any ${}_A H_{L(\mathbf{R})^T}$ sub-bimodule of ${}_A L^2(M^T)_{L(\mathbf{R})^T}$ such that $\dim(H_{L(\mathbf{R})^T}) < \infty$, one has $H \subset L^2(L(\mathbf{R})^T)$. In particular $A' \cap M^T \subset L(\mathbf{R})^T$.

Corollary 3.8. *Let $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ be a free Araki-Woods factor such that the orthogonal representation (U_t) is mixing. Then the continuous core $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ is a solid II_{∞} factor.*

Proof of Theorem 3.7. As usual, denote by (λ_t) the unitaries in $L(\mathbf{R})$ that implement the modular action σ on $\mathcal{M}$. Let $\Phi : L^{\infty}(\mathbf{R}) \rightarrow L(\mathbf{R})$ be the Fourier Transform so that $\Phi(e^{it \cdot}) = \lambda_t$, for every $t \in \mathbf{R}$. Let $T > 0$ and denote by $q = \Phi(\chi_{[0,T]})$. Notice that $L^{\infty}(\mathbf{R})\chi_{[0,T]} \cong L^{\infty}[0, T]$ and that

$$\text{span} \left\{ \sum_{k \in F} c_k e^{i \frac{2\pi}{T} k \cdot} \chi_{[0,T]} : F \subset \mathbf{Z} \text{ finite subset, } c_k \in \mathbf{C}, \forall k \in F \right\}$$

is a unital $*$ -strongly dense $*$ -subalgebra of $L^{\infty}(\mathbf{R})\chi_{[0,T]}$. Thus, using the isomorphism Φ , we get that

$$\mathcal{A} := \text{span} \left\{ \sum_{k \in F} c_k \lambda_{\frac{2\pi}{T} k} q : F \subset \mathbf{Z} \text{ finite subset, } c_k \in \mathbf{C}, \forall k \in F \right\}$$

is a unital $*$ -strongly dense $*$ -subalgebra of $L(\mathbf{R})q$. Let (u_n) be bounded sequence in $L(\mathbf{R})q$ such that $u_n \rightarrow 0$ weakly, as $n \rightarrow \infty$, and $\|u_n\|_{\infty} \leq 1$, for every $n \in \mathbf{N}$. Using Kaplansky density theorem together with a standard diagonal process, choose a sequence $y_n \in \mathcal{A}$ such that $\|y_n\|_{\infty} \leq 1$, for every $n \in \mathbf{N}$, and $\|u_n - y_n\|_{2, \text{Tr}} \rightarrow 0$, as $n \rightarrow \infty$. We will write $y_n = z_n q$ with

$$z_n = \sum_{k \in F_n} c_{k,n} \lambda_{\frac{2\pi}{T} k},$$

where $F_n \subset \mathbf{Z}$ is finite, $c_{k,n} \in \mathbf{C}$, for any $k \in F_n$ and any $n \in \mathbf{N}$. Using the T -periodicity, we have for any $n \in \mathbf{N}$,

$$\begin{aligned} \|z_n\|_{\infty} &= \|\Phi^{-1}(z_n)\|_{\infty} \\ &= \text{ess sup}_{x \in \mathbf{R}} \left| \sum_{k \in F_n} c_{k,n} e^{i \frac{2\pi}{T} k x} \right| \\ &= \text{ess sup}_{x \in [0, T]} \left| \sum_{k \in F_n} c_{k,n} e^{i \frac{2\pi}{T} k x} \right| \\ &= \|\Phi^{-1}(z_n)\chi_{[0,T]}\|_{\infty} \\ &= \|y_n\|_{\infty} \leq 1. \end{aligned}$$

Thus, the sequence (z_n) is uniformly bounded.

The **first step** of the proof consists in proving the following:

$$\|E_{L(\mathbf{R})q}(au_n b)\|_{2, \text{Tr}} \rightarrow 0, \text{ as } n \rightarrow \infty, \forall a, b \in qMq \cap \ker(E_{L(\mathbf{R})q}).$$

Equivalently, we need to show that

$$(7) \quad \|qE_{L(\mathbf{R})}(au_n b)q\|_{2, \text{Tr}} \rightarrow 0, \text{ as } n \rightarrow \infty, \forall a, b \in \ker(E_{L(\mathbf{R})}).$$

The **first step** of the proof is now divided in three different claims that will lead to proving (7). First note that

$$\mathcal{E} := \text{span} \left\{ \sum_{t \in F} x_t \lambda_t : F \subset \mathbf{R} \text{ finite subset, } x_t \in \mathcal{M} \text{ with } \varphi(x_t) = 0, \forall t \in F \right\}$$

is $*$ -strongly dense in $\ker(E_L(\mathbf{R}))$ by Kaplansky density theorem. We first prove the following:

Claim 3.9. *If $\|qE_L(\mathbf{R})(xu_ny)q\|_{2,\text{Tr}} \rightarrow 0$, as $n \rightarrow \infty$, $\forall x, y \in \mathcal{M}$ with $\varphi(x) = \varphi(y) = 0$, then (7) is satisfied.*

Proof of Claim 3.9. Assume $\|qE_L(\mathbf{R})(xu_ny)q\|_{2,\text{Tr}} \rightarrow 0$, as $n \rightarrow \infty$, $\forall x, y \in \mathcal{M}$ with $\varphi(x) = \varphi(y) = 0$. First take $a \in \mathcal{E}$ that we write $a = \sum_{s \in F} x_s \lambda_s$, with $F \subset \mathbf{R}$ finite subset, such that $x_s \in \mathcal{M}$, $\varphi(x_s) = 0$, for every $s \in F$. Then take $b \in \ker(E_L(\mathbf{R}))$ and let $(b_j)_{j \in J}$ be a sequence in $\mathcal{E}$ such that $b - b_j \rightarrow 0$ $*$ -strongly, as $j \rightarrow \infty$. Since $\|u_n\|_\infty \leq 1$, we get for any $n \in \mathbf{N}$ and any $j \in J$,

$$\begin{aligned} \|qE_L(\mathbf{R})(au_nb)q\|_{2,\text{Tr}} &\leq \|qE_L(\mathbf{R})(au_nb_j)q\|_{2,\text{Tr}} + \|qE_L(\mathbf{R})(au_n(b - b_j))q\|_{2,\text{Tr}} \\ &\leq \|qE_L(\mathbf{R})(au_nb_j)q\|_{2,\text{Tr}} + \|a\|_\infty \|(b - b_j)q\|_{2,\text{Tr}} \end{aligned}$$

Fix $\varepsilon > 0$. Since $b - b_j \rightarrow 0$ $*$ -strongly, as $j \rightarrow \infty$, fix $j_0 \in J$ such that $\|(b - b_{j_0})q\|_{2,\text{Tr}} \leq \varepsilon/2$. Write $b_{j_0} = \sum_{t \in F'} y_t \lambda_t$, with $F' \subset \mathbf{R}$ finite subset, such that $y_t \in \mathcal{M}$, $\varphi(y_t) = 0$, for every $t \in F'$. Therefore, for any $n \in \mathbf{N}$,

$$\begin{aligned} \|qE_L(\mathbf{R})(au_nb_{j_0})q\|_{2,\text{Tr}} &\leq \sum_{(s,t) \in F \times F'} \|qE_L(\mathbf{R})(x_s \lambda_s u_n y_t \lambda_t)q\|_{2,\text{Tr}} \\ &= \sum_{(s,t) \in F \times F'} \|\lambda_s qE_L(\mathbf{R})(\sigma_{-s}(x_s) u_n y_t) q \lambda_t\|_{2,\text{Tr}} \\ &= \sum_{(s,t) \in F \times F'} \|qE_L(\mathbf{R})(\sigma_{-s}(x_s) u_n y_t) q\|_{2,\text{Tr}}. \end{aligned}$$

Since $\varphi(\sigma_{-s}(x_s)) = \varphi(y_t) = 0$, for any $(s, t) \in F \times F'$, using the assumption of the claim, there exists $n_0 \in \mathbf{N}$ large enough such that for any $n \geq n_0$, $\|qE_L(\mathbf{R})(au_nb_{j_0})q\|_{2,\text{Tr}} \leq \varepsilon/2$. Thus, for any $n \geq n_0$, $\|qE_L(\mathbf{R})(au_nb)q\|_{2,\text{Tr}} \leq \varepsilon$. This proves that for any $a \in \mathcal{E}$ and any $b \in \ker(E_L(\mathbf{R}))$, $\|qE_L(\mathbf{R})(au_nb)q\|_{2,\text{Tr}} \rightarrow 0$, as $n \rightarrow \infty$. If we do the same thing by approximating $a \in \ker(E_L(\mathbf{R}))$ with elements in $\mathcal{E}$, we finally get the claim. $\square$

We now replace the sequence (u_n) by (z_n) , use the mixing property of the modular action σ and prove the following:

Claim 3.10. $\forall a, b \in (\mathcal{M})_1$ with $\varphi(a) = \varphi(b) = 0$, $\|qE_L(\mathbf{R})(az_nb)q\|_{2,\text{Tr}} \rightarrow 0$, as $n \rightarrow \infty$.

Proof of Claim 3.10. Fix $a, b \in (\mathcal{M})_1$ such that $\varphi(a) = \varphi(b) = 0$. Fix $\varepsilon > 0$. For any $n \in \mathbf{N}$, we have

$$\begin{aligned} \|qE_L(\mathbf{R})(az_nb)q\|_{2,\text{Tr}}^2 &= \left\| \sum_{k \in F_n} c_k \varphi\left(a \sigma_{\frac{2\pi}{T}k}(b)\right) \lambda_k q \right\|_{2,\text{Tr}}^2 \\ &= \text{Tr}(q) \sum_{k \in F_n} |c_k|^2 \left| \varphi\left(a \sigma_{\frac{2\pi}{T}k}(b)\right) \right|^2. \end{aligned}$$

Moreover for any $n \in \mathbf{N}$,

$$\mathrm{Tr}(q) \sum_{k \in F_n} |c_k|^2 = \|z_n q\|_{2, \mathrm{Tr}}^2 \leq \mathrm{Tr}(q) \|z_n q\|_{\infty}^2 \leq T.$$

Since the modular group σ is φ -mixing (because (U_t) is assumed to be mixing), there exists a finite subset $K \subset \mathbf{Z}$ such that for any $k \in \mathbf{Z} \setminus K$, $\left| \varphi \left(a \sigma_{\frac{2\pi}{T}k}(b) \right) \right| \leq \varepsilon / \sqrt{2T}$. Thus,

$$\|q E_{L(\mathbf{R})}(a z_n b) q\|_{2, \mathrm{Tr}} \leq \left\| \sum_{k \in K \cap F_n} c_k \lambda_{\frac{2\pi}{T}k} q \right\|_{2, \mathrm{Tr}} + \varepsilon/2.$$

Since $u_n - z_n q \rightarrow 0$ strongly and $u_n \rightarrow 0$ weakly, as $n \rightarrow \infty$, it follows that $z_n q \rightarrow 0$ weakly, as $n \rightarrow \infty$. In particular there exists n_0 large enough such that for any $n \geq n_0$, for any $k \in K \cap F_n$, $|c_k| \leq \varepsilon / (2|K| \|q\|_{2, \mathrm{Tr}})$. Thus, for any $n \geq n_0$,

$$\|q E_{L(\mathbf{R})}(a z_n b) q\|_{2, \mathrm{Tr}} \leq \varepsilon/2 + \varepsilon/2 = \varepsilon.$$

This proves that $\|q E_{L(\mathbf{R})}(a z_n b) q\|_{2, \mathrm{Tr}} \rightarrow 0$, as $n \rightarrow \infty$. $\square$

The last claim consists in going back to the sequence (u_n) and proving the following:

Claim 3.11. $\forall a, b \in (\mathcal{M})_1$ with $\varphi(a) = \varphi(b) = 0$, $\|q E_{L(\mathbf{R})}(a u_n b) q\|_{2, \mathrm{Tr}} \rightarrow 0$, as $n \rightarrow \infty$.

Proof of Claim 3.11. Applying once more Kaplansky density theorem, we can find a sequence $(q_i)_{i \in I}$ in $L(\mathbf{R})$ such that

- $q_i = \sum_{t \in F_i} d_t \lambda_t$, with $F_i \subset \mathbf{R}$ finite subset, $d_t \in \mathbf{C}$, for any $t \in F_i$ and for any $i \in I$;
- $\|q_i\|_{\infty} \leq 1$, for any $i \in I$;
- $q - q_i \rightarrow 0$ *-strongly, as $i \rightarrow \infty$.

Fix now $a, b \in (\mathcal{M})_1$ such that $\varphi(a) = \varphi(b) = 0$. Using the fact that

$$\|a\|_{\infty}, \|b\|_{\infty}, \|q\|_{\infty}, \|z_n\|_{\infty} \leq 1, \forall n \in \mathbf{N},$$

we get for any $n \in \mathbf{N}$ and any $i \in I$,

$$\begin{aligned} \|q E_{L(\mathbf{R})}(a u_n b) q\|_{2, \mathrm{Tr}} &\leq \|q E_{L(\mathbf{R})}(a(u_n - z_n q)b) q\|_{2, \mathrm{Tr}} + \|q E_{L(\mathbf{R})}(a z_n q b) q\|_{2, \mathrm{Tr}} \\ &\leq \|u_n - z_n q\|_{2, \mathrm{Tr}} + \|q E_{L(\mathbf{R})}(a z_n (q - q_i)b) q\|_{2, \mathrm{Tr}} \\ &\quad + \|q E_{L(\mathbf{R})}(a z_n q_i b) q\|_{2, \mathrm{Tr}} \\ &\leq \|u_n - z_n q\|_{2, \mathrm{Tr}} + \|(q - q_i) b q\|_{2, \mathrm{Tr}} \\ &\quad + \sum_{t \in F_i} |d_t| \|q E_{L(\mathbf{R})}(a z_n \sigma_t(b)) \lambda_t q\|_{2, \mathrm{Tr}} \\ &\leq \|u_n - z_n q\|_{2, \mathrm{Tr}} + \|(q - q_i) b q\|_{2, \mathrm{Tr}} \\ &\quad + \sum_{t \in F_i} |d_t| \|q E_{L(\mathbf{R})}(a z_n \sigma_t(b)) q\|_{2, \mathrm{Tr}}. \end{aligned}$$

Since $q - q_i \rightarrow 0$ *-strongly, as $i \rightarrow \infty$, it follows that $\|(q - q_i) b q\|_{2, \mathrm{Tr}} \rightarrow 0$, as $i \rightarrow \infty$. Fix $\varepsilon > 0$. Then, take $i_0 \in I$ such that $\|(q - q_{i_0}) b q\|_{2, \mathrm{Tr}} \leq \varepsilon/3$. Since $\|u_n - z_n q\|_{2, \mathrm{Tr}} \rightarrow 0$, as $n \rightarrow \infty$ and using Claim 3.10, we may choose n_0 large enough such that for any $n \geq n_0$,

$$\begin{aligned} \|u_n - z_n q\|_{2, \mathrm{Tr}} &\leq \varepsilon/3 \\ \sum_{t \in F_{i_0}} |d_t| \|q E_{L(\mathbf{R})}(a z_n \sigma_t(b)) q\|_{2, \mathrm{Tr}} &\leq \varepsilon/3. \end{aligned}$$

Consequently, for any $n \geq n_0$, we get $\|qE_{L(\mathbf{R})}(au_nb)q\|_{2,\text{Tr}} \leq \varepsilon$. Therefore, we have proven $\|qE_{L(\mathbf{R})}(au_nb)q\|_{2,\text{Tr}} \rightarrow 0$, as $n \rightarrow \infty$. $\square$

Thanks to Claims 3.9 and 3.11, it is then clear that (7) is satisfied. This finishes the **first step** of the proof.

The **last step** of the proof consists in using Theorem 2.8. Let $k \geq 1$ and $q \in \mathbf{M}_k(\mathbf{C}) \otimes L(\mathbf{R})$ be a non-zero projection such that $T := (\text{Tr}_k \otimes \text{Tr})(q) < \infty$. Since $\mathbf{M}_k(\mathbf{C}) \otimes M$ is a II_∞ factor, there exists a unitary $u \in \mathcal{U}(\mathbf{M}_k(\mathbf{C}) \otimes M)$ such that

$$q = u \begin{pmatrix} q_0 & & 0 \\ & \ddots & \\ 0 & & q_0 \end{pmatrix} u^*$$

where $q_0 = \Phi(\chi_{[0,T/k]}) \in L(\mathbf{R})$. Using the spatiality of $\text{Ad}(u)$ on $\mathbf{M}_k(\mathbf{C}) \otimes M$, we may assume without loss of generality that

$$q = \begin{pmatrix} q_0 & & 0 \\ & \ddots & \\ 0 & & q_0 \end{pmatrix}$$

In particular, $q \in \mathbf{M}_k(\mathbf{C}) \otimes L(\mathbf{R})q_0$. Define $M^T := q(\mathbf{M}_k(\mathbf{C}) \otimes M)q$ and $L(\mathbf{R})^T := q(\mathbf{M}_k(\mathbf{C}) \otimes L(\mathbf{R}))q$. Let $A \subset L(\mathbf{R})^T$ be a diffuse von Neumann subalgebra. Choose a sequence of unitaries (u_n) in A such that $u_n \rightarrow 0$ weakly, as $n \rightarrow \infty$. Thus, we can write $u_n = [u_n^{i,j}]_{i,j}$ where $u_n^{i,j} \in L(\mathbf{R})q_0$ and $\|u_n^{i,j}\|_\infty \leq 1$, for any $n \in \mathbf{N}$ and any $i, j \in \{1, \dots, k\}$. Moreover, $u_n^{i,j} \rightarrow 0$ weakly, as $n \rightarrow \infty$, in $L(\mathbf{R})q_0$, for any $i, j \in \{1, \dots, k\}$. Thus, using the **first step** of the proof, it becomes clear that the inclusion $L(\mathbf{R})^T \subset M^T$ is weakly mixing through A in the sense of Definition 2.7. Thus, using Theorem 2.8, it follows that for any ${}_A H_{L(\mathbf{R})^T}$ sub-bimodule of ${}_A L^2(M^T)_{L(\mathbf{R})^T}$ such that $\dim(H_{L(\mathbf{R})^T}) < \infty$, one has $H \subset L^2(L(\mathbf{R})^T)$. In particular $A' \cap M^T \subset L(\mathbf{R})^T$. $\square$

Proof of Corollary 3.8. Let $q \in L(\mathbf{R})$ be a non-zero projection such that $\text{Tr}(q) < \infty$. Denote by $N = qMq$ the corresponding II_1 factor. By contradiction assume that N is not solid. Then there exists a non-amenable von Neumann subalgebra $Q \subset N$ such that the relative commutant $Q' \cap N$ is diffuse. Since N is a II_1 factor, using the same argument as in the proof of Corollary 3.5, we may assume that Q has no amenable direct summand and $Q_0 = Q' \cap N$ is still diffuse.

Since Q has no amenable direct summand, Theorem 3.4 yields $Q_0 \preceq_M L(\mathbf{R})$. Thus using Theorem 2.9, we know that there exists a non-zero projection $p \in L(\mathbf{R})$ such that $\text{Tr}(p) < \infty$, and $Q_0 \preceq_{eMe} L(\mathbf{R})p$ where $e = p \vee q$. Consequently, there exist $n \geq 1$, a (possibly non-unital) $*$ -homomorphism $\psi : Q_0 \rightarrow \mathbf{M}_n(\mathbf{C}) \otimes L(\mathbf{R})p$ and a non-zero partial isometry $v \in \mathbf{M}_{1,n}(\mathbf{C}) \otimes qMp$ such that

$$xv = v\psi(x), \forall x \in Q_0.$$

We moreover have

$$vv^* \in Q'_0 \cap qMq \text{ and } v^*v \in \psi(Q'_0)' \cap \psi(q)(\mathbf{M}_n(\mathbf{C}) \otimes pMp)\psi(q).$$

Write $Q_1 = Q'_0 \cap qMq$ and notice that $Q \subset Q_1$. Since $\psi(Q_0)$ is diffuse and $v^*v \in \psi(Q'_0)' \cap \psi(q)(\mathbf{M}_n(\mathbf{C}) \otimes pMp)\psi(q)$, Theorem 3.7 yields $v^*v \in \psi(q)(\mathbf{M}_n(\mathbf{C}) \otimes L(\mathbf{R})p)\psi(q)$, so that we may assume $v^*v = \psi(q)$. For any $y \in Q_1$, and any $x \in Q_0$,

$$\begin{aligned} v^*yv\psi(x) &= v^*yxv \\ &= v^*xyv \\ &= \psi(x)v^*yv. \end{aligned}$$

Thus, $v^*Q_1v \subset \psi(Q_0)' \cap v^*v(\mathbf{M}_n(\mathbf{C}) \otimes pMp)v^*v$. Since $\psi(Q_0)$ is diffuse, Theorem 3.7 yields $v^*Q_1v \subset v^*v(\mathbf{M}_n(\mathbf{C}) \otimes L(\mathbf{R})p)v^*v$. Since Q has no amenable direct summand and $Q \subset Q_1$ is a unital von Neumann subalgebra, it follows that Q_1 has no amenable direct summand either. Thus the von Neumann algebra $vv^*Q_1vv^*$ is non-amenable. But $\text{Ad}(v^*) : vv^*Mvv^* \rightarrow v^*v(\mathbf{M}_n(\mathbf{C}) \otimes pMp)v^*v$ is a $*$ -isomorphism and

$$\text{Ad}(v^*)(vv^*Q_1vv^*) \subset v^*v(\mathbf{M}_n(\mathbf{C}) \otimes L(\mathbf{R})p)v^*v.$$

Since $v^*v(\mathbf{M}_n(\mathbf{C}) \otimes L(\mathbf{R})p)v^*v$ is of type I, hence amenable, we get a contradiction. $\square$

Since the left regular representation (λ_t) of $\mathbf{R}$ acting on $L^2_{\mathbf{R}}(\mathbf{R}, \text{Lebesgue})$ is mixing, the continuous core M of $\Gamma(L^2_{\mathbf{R}}(\mathbf{R}, \text{Lebesgue}), \lambda_t)''$ is solid. We partially retrieve a previous result of Shlyakhtenko [31] where he proved in this case that $M \cong L(\mathbf{F}_{\infty}) \otimes \mathbf{B}(\ell^2)$, which is solid by [15]. We will give in the next section an example of a non-amenable solid II_1 factor with full fundamental group which is not isomorphic to any interpolated free group factor $L(\mathbf{F}_t)$, for $1 < t \leq \infty$.

Note that the mixing property of the representation (U_t) is not a necessary condition for the solidity of the continuous core M . Indeed, take $U_t = \text{Id} \oplus \lambda_t$ on $H_{\mathbf{R}} = \mathbf{R} \oplus L^2_{\mathbf{R}}(\mathbf{R}, \text{Lebesgue})$. Then (U_t) is not mixing, but the continuous core M of $\Gamma(H_{\mathbf{R}}, U_t)''$ is still isomorphic to $L(\mathbf{F}_{\infty}) \otimes \mathbf{B}(\ell^2)$ [29].

4. EXAMPLES OF SOLID II_1 FACTORS

4.1. Probability measures on the real line and unitary representations of $\mathbf{R}$. Write λ for the Lebesgue measure on the real line $\mathbf{R}$. Let μ be a *symmetric* (positive) probability measure on $\mathbf{R}$, i.e. $\mu(X) = \mu(-X)$, for any Borel subset $X \subset \mathbf{R}$. Consider the following unitary representation (U_t^{μ}) of $\mathbf{R}$ on $L^2(\mathbf{R}, \mu)$ given by:

$$(8) \quad (U_t^{\mu}f)(x) = e^{itx}f(x), \forall f \in L^2(\mathbf{R}, \mu), \forall t, x \in \mathbf{R}.$$

Define the Hilbert subspace of $L^2(\mathbf{R}, \mu)$

$$(9) \quad K_{\mathbf{R}}^{\mu} := \left\{ f \in L^2(\mathbf{R}, \mu) : f(x) = \overline{f(-x)}, \forall x \in \mathbf{R} \right\}.$$

Since μ is assumed to be symmetric, the restriction of the inner product to $K_{\mathbf{R}}^{\mu}$ is real-valued. Indeed, for any $f, g \in K_{\mathbf{R}}^{\mu}$,

$$\begin{aligned} \langle f, g \rangle &= \int_{\mathbf{R}} f(x) \overline{g(x)} d\mu(x) \\ &= \int_{\mathbf{R}} f(-x) \overline{g(-x)} d\mu(-x) \\ &= \int_{\mathbf{R}} \overline{f(x)} g(x) d\mu(x) \\ &= \overline{\langle f, g \rangle}. \end{aligned}$$

Moreover the representation (U_t^{μ}) leaves $K_{\mathbf{R}}^{\mu}$ globally invariant. Thus, (U_t^{μ}) restricted to $K_{\mathbf{R}}^{\mu}$ becomes an orthogonal representation. Define the *Fourier Transform* of the probability measure μ by:

$$\tilde{\mu}(t) = \int_{\mathbf{R}} e^{itx} d\mu(x), \forall t \in \mathbf{R}.$$

We shall identify $\widehat{\mathbf{R}}$ with $\mathbf{R}$ in the usual way, such that

$$\widehat{f}(t) = \int_{\mathbf{R}} e^{itx} f(x) d\lambda(x), \forall t \in \mathbf{R}, \forall f \in L^1(\mathbf{R}, \lambda).$$

Proposition 4.1. *Let μ be a symmetric probability measure on $\mathbf{R}$. Then*

$$(U_t^\mu) \text{ is mixing} \iff \tilde{\mu}(t) \rightarrow 0, \text{ as } |t| \rightarrow \infty.$$

Proof. We prove both directions.

$\implies$ Assume (U_t^μ) is mixing. Let $f = \mathbf{1}_{\mathbf{R}} \in L^2(\mathbf{R}, \mu)$ be the constant function equal to 1. Then

$$\begin{aligned} \tilde{\mu}(t) &= \int_{\mathbf{R}} e^{itx} d\mu(x) \\ &= \langle U_t^\mu f, f \rangle \rightarrow 0, \text{ as } |t| \rightarrow \infty. \end{aligned}$$

$\impliedby$ Assume $\tilde{\mu}(t) \rightarrow 0$, as $|t| \rightarrow \infty$. Let $f, g \in L^2(\mathbf{R}, \mu)$. Then $h := f\bar{g} \in L^1(\mathbf{R}, \mu)$. Since the set $\{f \in C_0(\mathbf{R}) : \hat{f} \in L^1(\mathbf{R}, \lambda)\}$ is dense in $L^1(\mathbf{R}, \mu)$, we may choose a sequence (h_n) in $C_0(\mathbf{R})$ such that $\|h - h_n\|_{L^1(\mathbf{R}, \mu)} \rightarrow 0$, as $n \rightarrow \infty$, and $\hat{h}_n \in L^1(\mathbf{R}, \lambda)$, for any $n \in \mathbf{N}$. Define

$$\begin{aligned} \tilde{h}(t) &= \int_{\mathbf{R}} e^{itx} h(x) d\mu(x), \forall t \in \mathbf{R} \\ \tilde{h}_n(t) &= \int_{\mathbf{R}} e^{itx} h_n(x) d\mu(x), \forall t \in \mathbf{R}, \forall n \in \mathbf{N}. \end{aligned}$$

Since $\|h - h_n\|_{L^1(\mathbf{R}, \mu)} \rightarrow 0$, as $n \rightarrow \infty$, it follows that $\|\tilde{h} - \tilde{h}_n\|_\infty \rightarrow 0$, as $n \rightarrow \infty$. Since $\hat{h}_n \in L^1(\mathbf{R}, \lambda)$, we know that

$$h_n(x) = C \int_{\mathbf{R}} e^{-ixu} \hat{h}_n(u) d\lambda(u), \forall x \in \mathbf{R},$$

where C is a universal constant that only depends on the normalization of the Lebesgue measure λ on $\mathbf{R}$. Therefore, for any $t \in \mathbf{R}$ and any $n \in \mathbf{N}$,

$$\begin{aligned} \tilde{h}_n(t) &= \int_{x \in \mathbf{R}} e^{itx} h_n(x) d\mu(x) \\ &= C \int_{x \in \mathbf{R}} \left(\int_{u \in \mathbf{R}} e^{i(t-u)x} \hat{h}_n(u) d\lambda(u) \right) d\mu(x) \\ &= C \int_{u \in \mathbf{R}} \hat{h}_n(u) \left(\int_{x \in \mathbf{R}} e^{i(t-u)x} d\mu(x) \right) d\lambda(u) \\ &= C \int_{u \in \mathbf{R}} \hat{h}_n(u) \tilde{\mu}(t-u) d\lambda(u) \\ &= C \left(\hat{h}_n * \tilde{\mu} \right)(t), \end{aligned}$$

where $*$ is the convolution product. Since $\tilde{\mu} \in C_0(\mathbf{R})$ and $\hat{h}_n \in L^1(\mathbf{R}, \lambda)$, it is easy to check that $\hat{h}_n * \tilde{\mu} \in C_0(\mathbf{R})$. Consequently, $\tilde{h}_n \in C_0(\mathbf{R})$ and since $\|\tilde{h} - \tilde{h}_n\|_\infty \rightarrow 0$, as $n \rightarrow \infty$, it follows that $\tilde{h} \in C_0(\mathbf{R})$. But for any $t \in \mathbf{R}$,

$$\begin{aligned} \langle U_t^\mu f, g \rangle &= \int_{\mathbf{R}} e^{itx} f(x) \overline{g(x)} d\mu(x) \\ &= \tilde{h}(t). \end{aligned}$$

Thus, the unitary representation (U_t^μ) is mixing. □

For a measure ν on $\mathbf{R}$, define the *measure class* of ν by:

$$\mathcal{C}_\nu := \{\nu' : \nu' \text{ is absolutely continuous w.r.t. } \nu\}.$$

Definition 4.2. Let (V_t) be a unitary representation of $\mathbf{R}$ on a separable Hilbert space H . Denote by B the infinitesimal generator of (V_t) , i.e. B is the positive, self-adjoint (possibly) unbounded operator on H such that $V_t = B^{it}$, for every $t \in \mathbf{R}$. We define the *spectral measure* of the representation (V_t) as the spectral measure of the operator B and denote it by $\mathcal{C}_V = \mathcal{C}_B$.

The *measure class* $\mathcal{C}_V$ can also be defined as the smallest collection of all the measures ν on $\mathbf{R}$ such that:

- (1) If $\nu \in \mathcal{C}_V$ and ν' is absolutely continuous w.r.t. ν , then $\nu' \in \mathcal{C}_V$;
- (2) For any unit vector $\eta \in H$, the probability measure associated with the positive definite function $t \mapsto \langle V_t \eta, \eta \rangle$ belongs to $\mathcal{C}_V$.

Since H is separable, there exists a measure ν that generates $\mathcal{C}_V$, i.e. $\mathcal{C}_V$ is the smallest collection of measures on $\mathbf{R}$ satisfying (1) and containing ν . We will refer to this particular measure ν as the “spectral measure” of the representation (V_t) and simply denote it by ν .

Let μ be a symmetric probability measure on $\mathbf{R}$ and consider the unitary representation (U_t^μ) on $L^2(\mathbf{R}, \mu)$ as defined in (8). Then for any unit vector $f \in L^2(\mathbf{R}, \mu)$,

$$\langle U_t^\mu f, f \rangle = \int_{\mathbf{R}} e^{itx} |f(x)|^2 d\mu(x), \forall t \in \mathbf{R}.$$

Since the probability measure $|f(x)|^2 d\mu(x)$ is absolutely continuous w.r.t. $d\mu(x)$, it is clear that the spectral measure of (U_t^μ) is μ . More generally, we have the following:

Proposition 4.3. *Let μ be a symmetric probability measure on $\mathbf{R}$. Consider the unitary representation (U_t^μ) defined on $L^2(\mathbf{R}, \mu)$ by (8). Then for any $n \geq 1$, the spectral measure of the n -fold tensor product $(U_t^\mu)^{\otimes n}$ is the n -fold convolution product*

$$\mu^{*n} = \underbrace{\mu * \cdots * \mu}_{n \text{ times}}.$$

4.2. Examples of solid II_1 factors. Erdős showed in [9] that the symmetric probability measure μ_θ , with $\theta = 5/2$, obtained as the weak limit of

$$\left(\frac{1}{2} \delta_{-\theta^{-1}} + \frac{1}{2} \delta_{\theta^{-1}} \right) * \cdots * \left(\frac{1}{2} \delta_{-\theta^{-n}} + \frac{1}{2} \delta_{\theta^{-n}} \right)$$

has a Fourier Transform

$$\tilde{\mu}_\theta(t) = \prod_{n \geq 1} \cos \left(\frac{t}{\theta^n} \right)$$

which vanishes at infinity, i.e. $\tilde{\mu}(t) \rightarrow 0$, as $|t| \rightarrow \infty$, and μ_θ is singular w.r.t. the Lebesgue measure λ .

Example 4.4. Modifying the measure μ_θ , Antoniou & Shkarin (see Theorem 2.5, v in [2]) constructed an example of a symmetric probability μ on $\mathbf{R}$ such that:

- (1) The Fourier Transform of μ vanishes at infinity, i.e. $\tilde{\mu}(t) \rightarrow 0$, as $|t| \rightarrow \infty$.
- (2) For any $n \geq 1$, the n -fold convolution product μ^{*n} is singular w.r.t. the Lebesgue measure λ .

Let μ be a symmetric probability measure on $\mathbf{R}$ as in Example 4.4. Proposition 4.1 and Proposition 4.3 yields that the unitary representation (U_t^μ) defined on $L^2(\mathbf{R}, \mu)$ by (8) satisfies:

- (1) (U_t^μ) is mixing.
- (2) The spectral measure of $\bigoplus_{n \geq 1} (U_t^\mu)^{\otimes n}$ is singular w.r.t. the Lebesgue measure λ .

Let now $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ and let $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ be the continuous core. Let $q \in L(\mathbf{R})$ be a non-zero projection such that $\text{Tr}(q) < \infty$. Denote by $N = qMq$ the corresponding II_1 factor. Using free probability techniques such as the *free entropy*, Shlyakhtenko (see Theorem 9.12 in [30]) showed that if the spectral measure of the unitary representation $\bigoplus_{n \geq 1} U_t^{\otimes n}$ is singular w.r.t. the Lebesgue measure λ , then for any finite set of generators $X_1, \dots, X_n$ of N , the free entropy dimension satisfies

$$\delta_0(X_1, \dots, X_n) \leq 1.$$

In particular, N is not isomorphic to any interpolated free group factor $L(\mathbf{F}_t)$, for $1 < t \leq \infty$. Combining these two results together with Corollary 3.7, we obtain the following:

Theorem 4.5. *Let μ be a symmetric probability measure on $\mathbf{R}$ as in Example 4.4. Let $\mathcal{M} = \Gamma(K_{\mathbf{R}}^{\mu}, U_t^{\mu})''$ be the free Araki-Woods factor associated with the orthogonal representation (U_t^{μ}) acting on the real Hilbert space $K_{\mathbf{R}}^{\mu}$, as defined in (8–9). Let $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ be the continuous core. Fix a non-zero projection $q \in L(\mathbf{R})$ such that $\text{Tr}(q) < \infty$, and denote by $N = qMq$ the corresponding II_1 factor. Then*

- (1) N is non-amenable and solid.
- (2) N has full fundamental group, i.e. $\mathcal{F}(N) = \mathbf{R}_+^*$.
- (3) N is not isomorphic to any interpolated free group factor $L(\mathbf{F}_t)$, for $1 < t \leq \infty$.

We believe that all the free Araki-Woods factors $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ have the *complete metric approximation property* (c.m.a.p.), i.e. there exists a sequence $\Phi_n : \mathcal{M} \rightarrow \mathcal{M}$ of finite rank, completely bounded maps such that $\Phi_n \rightarrow \text{Id}$ ultraweakly pointwise, as $n \rightarrow \infty$, and $\limsup_{n \rightarrow \infty} \|\Phi_n\|_{\text{cb}} \leq 1$. If $\mathcal{M} = \Gamma(H_{\mathbf{R}}, U_t)''$ had the c.m.a.p. then by [1], the continuous core $M = \mathcal{M} \rtimes_{\sigma} \mathbf{R}$ would have the c.m.a.p., as well as the II_1 factor qMq , for $q \in M$ non-zero finite projection. On the other hand, the wreath product II_1 factors $L(\mathbf{Z} \wr \mathbf{F}_n)$ do not have the c.m.a.p., for any $2 \leq n \leq \infty$, by [18]. Thus, we conjecture that the solid II_1 factors constructed in Theorem 4.5 are not isomorphic to $L(\mathbf{Z} \wr \mathbf{F}_n)$, for any $2 \leq n \leq \infty$.

REFERENCES

- [1] C. ANANTHARAMAN-DELAROCHE, *Amenable correspondences and approximation properties for von Neumann algebras*. Pacific J. Math. **171** (1995), 309–341.
- [2] I. ANTONIOU & S. A. SHKARIN, *Decay measures on locally compact abelian topological groups*. Proc. Roy. Soc. Edinburgh **131** (2001), 1257–1273.
- [3] L. BARNETT, *Free product von Neumann algebras of type III*. Proc. Amer. Math. Soc. **123** (1995), 543–553.
- [4] I. CHIFAN & A. IOANA, *Ergodic subequivalence relations induced by a Bernoulli action*. [arXiv:0802.2353](#)
- [5] I. CHIFAN & C. HOUDAYER, *Prime factors and amalgamated free products*. [arXiv:0805.1566](#)
- [6] A. CONNES, *Almost periodic states and factors of type III₁*. J. Funct. Anal. **16** (1974), 415–445.
- [7] A. CONNES, *Une classification des facteurs de type III*. Ann. Sci. École Norm. Sup. **6** (1973), 133–252.
- [8] K. DYKEMA, *Interpolated free group factors*. Pacific J. Math. **163** (1994), 123–135.
- [9] P. ERDÖS, *On a family of symmetric Bernoulli convolutions*. Amer. J. Math. **61** (1939), 974–976.
- [10] U. HAAGERUP, *Connes’ bicentralizer problem and uniqueness of the injective factor of type III₁*. Acta Math. **69** (1986), 95–148.
- [11] C. HOUDAYER, *Free Araki-Woods factors and Connes’ bicentralizer problem*. [arXiv:0809.3827](#)
- [12] C. HOUDAYER, *Construction of type II₁ factors with prescribed countable fundamental group*. J. reine angew Math., to appear. [arXiv:0704.3502](#)
- [13] C. HOUDAYER, *On some free products of von Neumann algebras which are free Araki-Woods factors*. Int. Math. Res. Notices. Vol. **2007**, article ID rnm098, 21 pages.
- [14] A. IOANA, J. PETERSON & S. POPA, *Amalgamated free products of w-rigid factors and calculation of their symmetry groups*. Acta Math. **200** (2008), 85–153.
- [15] N. OZAWA, *Solid von Neumann algebras*. Acta Math. **192** (2004), 111–117.
- [16] N. OZAWA, *A Kurosh-type theorem for type II₁ factors*. Int. Math. Res. Notices. Vol. 2006 : article ID 97560, 21 pages.
- [17] N. OZAWA, *An example of a solid von Neumann algebra*. [arXiv:0804.0288](#)

- [18] N. OZAWA & S. POPA, *On a class of II_1 factors with at most one Cartan subalgebra*. To appear in Ann. of Math. [arXiv:0706.3623](#)
- [19] S. POPA, *On the superrigidity of malleable actions with spectral gap*. J. Amer. Math. Soc. **21** (2008), 981–1000.
- [20] S. POPA, *On Ozawa’s property for free group factors*. Int. Math. Res. Notices. Vol. 2007 : article ID rnm036, 10 pages.
- [21] S. POPA, *Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups I*. Invent. Math. **165** (2006), 369–408.
- [22] S. POPA, *On a class of type II_1 factors with Betti numbers invariants*. Ann. of Math. **163** (2006), 809–899.
- [23] S. POPA, *Some rigidity results for non-commutative Bernoulli Shifts*. J. Funct. Anal. **230** (2006), 273–328.
- [24] S. POPA & S. VAES, *Strong rigidity of generalized Bernoulli actions and computations of their symmetry groups*. Adv. Math. **217** (2008), 833–872.
- [25] F. RĂDULESCU, *Random matrices, amalgamated free products and subfactors of the von Neumann algebra of a free group, of noninteger index*. Invent. Math. **115** (1994), 347–389.
- [26] F. RĂDULESCU, *A one-parameter group of automorphisms of $L(\mathbf{F}_\infty) \otimes \mathbf{B}(H)$ scaling the trace*. C. R. Acad. Sci. Paris Sr. I Math. **314** (1992), 1027–1032.
- [27] D. SHLYAKHTENKO, *Some estimates for non-microstates free entropy dimension, with applications to q -semicircular families*. Int. Math. Res. Notices **51** (2004), 2757–2772.
- [28] D. SHLYAKHTENKO, *On the classification of full factors of type III*. Trans. Amer. Math. Soc. **356** (2004), 4143–4159.
- [29] D. SHLYAKHTENKO, *On multiplicity and free absorption for free Araki-Woods factors*. [math.OA/0302217](#)
- [30] D. SHLYAKHTENKO, *A -valued semicircular systems*. J. Funct. Anal. **166** (1999), 1–47.
- [31] D. SHLYAKHTENKO, *Some applications of freeness with amalgamation*. J. reine angew. Math. **500** (1998), 191–212.
- [32] D. SHLYAKHTENKO, *Free quasi-free states*. Pacific J. Math. **177** (1997), 329–368.
- [33] M. TAKESAKI, *Duality for crossed products and structure of von Neumann algebras of type III*. Acta Math. **131** (1973), 249–310.
- [34] Y. UEDA, *Amalgamated free products over Cartan subalgebra*. Pacific J. Math. **191** (1999), 359–392.
- [35] S. VAES & R. VERGNIOUX, *The boundary of universal discrete quantum groups, exactness and factoriality*. Duke Math. J. **140** (2007), 35–84.
- [36] S. VAES, *Rigidity results for Bernoulli actions and their von Neumann algebras (after S. Popa)*. Séminaire Bourbaki, exposé 961. Astérisque **311** (2007), 237–294.
- [37] S. VAES, *États quasi-libres libres et facteurs de type III (d’après D. Shlyakhtenko)*. Séminaire Bourbaki, exposé 937, Astérisque **299** (2005), 329–350.
- [38] D.-V. VOICULESCU, K.J. DYKEMA & A. NICA, *Free random variables*. CRM Monograph Series **1**. American Mathematical Society, Providence, RI, 1992.

MATH DEPT, UCLA, 520 PORTOLA PLAZA, LA, CA 90095

E-mail address: cyril@math.ucla.edu